Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab (Jan-Dec 2008)

Percent Susceptibility of Organisms Isolated From Stool (Jan-Dec 2008)

Organism isolated from stool	TOTAL ISOLATION	Ampicillin	Tetracycline	Trimethoprim/Sulfamethoxazole	Chloramphenicol	Norfloxacin	Ciprofloxacin	Ofloxacin	Oxacillin	Clindamycin	Erythromycin	Gentamicin	Vancomycin	Meropenem
Salmonella Typhi	1	-	-	100 (1)	-	100 (1)	-	-	-	-	-	-	-	-
Salmonella Non-typhoidal	836	51 (766)	-	85 (820)	-	100 (818)	96 (66)	99 (194)	-	-	-	-	-	4 (100)
Shigella dysenteriae	2	50 (2)	-	100 (2)	-	100 (2)	-	-	-	-	-	-	-	-
Shigella flexneri	19	33 (18)	-	33 (18)	-	94 (18)	-	-	-	-	-	-	-	-
Shigella sonnei	83	80 (81)	-	5 (83)	-	100 (82)	100 (1)	-	-	-	-	-	-	-
Staphylococcus aureus	50	-	100 (1)	88 (48)	100 (6)	88 (8)	-	-	91 (45)	83 (29)	78 (27)	85 (27)	100 (49)	-
Vibrio cholerae O1 Inaba	24	87 (23)	-	0 (22)	81 (16)	-	-	-	-	-	-	-	-	-
Vibrio cholerae O1 Ogawa	13	62 (13)	55 (11)	64 (11)	100 (11)	-	-	-	-	-	-	-	-	-
Vibrio cholerae non O1, non O139	30	86 (28)	88 (8)	90 (29)	100 (4)	-	-	-	-	-	-	-	-	-