

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 44 hospitals , Jan - Dec 2009.

Organism isolated from stool	TOTAL ISOLATION	BETA - LACTAM			QUINOLONE			AMINOGLYCOSIDE	MISCELLANEOUS				
		AMPICILLIN	OXACILLIN	CEFOXITIN	NALIDIXIC ACID	CIPROFLOXACIN	NORFLOXACIN		GENTAMICIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE
<i>Salmonella</i> Typhi	43	61.5 ⁰⁰ (878)	-	-	-	-	94.7 ⁰⁰ (38)	-	-	-	-	62.5 ⁰⁰ (40)	-
<i>Salmonella</i> Non-typhoidal	904	47.7 (878)	-	-	72.7 (44)	28.6 (42)	99.8 (876)	87.3 (181)	-	0 (33)	77.2 (430)	83.6 (876)	48.5 (443)
<i>Shigella dysenteriae</i>	53	41.3 ⁰⁵ (46)	-	-	-	-	98 ⁰⁵ (50)	-	-	-	82.9 ⁰⁰ (35)	56.2 ⁰⁵ (48)	60 ⁰⁵ (35)
<i>Shigella flexneri</i>	27	18.5 (27)	-	-	-	50.0 (4)	88.9 (27)	100 (1)	-	-	28.6 (7)	22.2 (27)	10.5 (19)
<i>Shigella sonnei</i>	93	57.1 (91)	-	-	-	90.6 ⁰⁵ (32)	97.8 (91)	-	-	-	-	5.4 (93)	-
<i>Staphylococcus aureus</i>	66	5.1 ⁰⁵ (39)	93.9 (33)	92.1 (38)	-	-	91 ⁰⁷ (36)	95.7 (47)	95.0 (40)	90.2 (41)	-	98.5 (66)	-
<i>Vibrio cholerae</i> O1 Inaba	34	81.8 ⁰⁵ (33)	-	-	-	-	97.1 ⁰⁵ (34)	-	-	-	99.1 ⁰⁰ (914)	21.9 ⁰⁵ (32)	97 ⁰⁵ (33)
<i>Vibrio cholerae</i> O1 Ogawa	626	66.4 ⁰⁷ (563)	-	-	-	-	99.7 ⁰⁷ (607)	-	-	2.1 ⁰⁷ (144)	99.6 ⁰⁷ (516)	4.6 ⁰⁷ (609)	4.4 ⁰⁷ (562)
<i>Vibrio cholerae</i> non O1, non O139	193	78.9 ⁰⁷ (190)	-	-	-	91.7 ⁰⁵ (36)	98.1 ⁰⁷ (104)	-	-	-	96.5 ⁰⁷ (143)	85.9 ⁰⁷ (192)	90 ⁰⁷ (150)
<i>Vibrio parahaemolyticus</i>	393	3.4 (355)	-	-	-	89 ⁰⁷ (73)	-	-	-	-	-	99.2 (366)	98.3 (301)

2000-2009



2003-2009



2004-2009



2005-2009



2007-2009



2008-2009

