

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab (Jan-Dec 2010)

Organism isolated from stool	TOTAL ISOLATION	BETA - LACTAM			QUINOLONE			AMINOGLYCOSIDE	MISCELLANEOUS				
		AMPICILLIN	OXACILLIN	CEFOXITIN	NALIDIXIC ACID	CIPROFLOXACIN	NORFLOXACIN		GENTAMICIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE
<i>Salmonella Typhi</i>	53	57.1 ⁰⁰ (49)	-	-	-	-	95.7 ⁰⁰ (47)	-	-	-	80.6 ⁰⁰ (36)	64.6 ⁰⁰ (48)	-
<i>Salmonella Non-typhoidal</i>	1,211	49.7 (1160)	-	-	76.7 (90)	47.6 (164)	100 (1185)	79.9 (159)	-	1.0 (101)	82.0 (495)	79.3 (1186)	52.4 (458)
<i>Shigella dysenteriae</i>	55	43.1 ⁰⁵ (51)	-	-	-	-	98.1 ⁰⁵ (51)	-	-	-	82.4 ⁰⁰ (34)	59.2 ⁰⁵ (49)	56.8 ⁰⁵ (37)
<i>Shigella flexneri</i>	69	24.2 (66)	-	-	-	86.7 ⁰⁰ (30)	100 (66)	93.3 ⁰⁰ (90)	-	-	20.2 ⁰⁵ (104)	20.6 (68)	25.8 (31)
<i>Shigella sonnei</i>	223	60.7 (211)	-	-	-	93.8 ⁰⁵ (42)	99.1 (214)	-	-	-	-	2.9 (172)	-
<i>Staphylococcus aureus</i>	56	4.0 ⁰⁰ (50)	88.6 ⁰⁸ (70)	78.4 (37)	-	-	86.7 ⁰⁸ (30)	74.3 (35)	87.2 ⁰⁸ (94)	87.1 (70)	-	85.5 (55)	-
<i>Vibrio cholerae O1 Inaba</i>	160	86.9 (153)	-	-	-	-	100 (118)	-	-	-	33.6 (137)	3.4 (147)	98.6 (147)
<i>Vibrio cholerae O1 Ogawa</i>	180	84.4 (179)	-	-	-	-	100 (153)	-	-	2.9 (136)	96.1 (152)	3.4 (175)	26.8 (138)
<i>Vibrio cholerae non O1, non O139</i>	161	80.4 (158)	-	-	-	97.4 ⁰⁵ (38)	100 (59)	-	-	-	94.5 (128)	85.4 (158)	91.8 (134)
<i>Vibrio parahaemolyticus</i>	1325	1.4 (1291)	-	-	-	97.1 (34)	-	-	-	-	-	99.5 (1280)	98 (1167)

