

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 44 hospitals , Jan - Dec 2011.

Organism isolated from stool	TOTAL ISOLATION	BETA - LACTAM	QUINOLONE			AMINOGLYCOSIDE	MISCELLANEOUS			
		AMPICILLIN	NALIDIXIC ACID	CIPROFLOXACIN	NORFLOXACIN	GENTAMICIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> Non-typhoidal	1125	43.2 (1044)	67.8 (90)	60.7 (183)	99.4 (1051)	64.7 (102)	2.1 (47)	77.1 (512)	84.3 (1090)	49.1 (589)
<i>Shigella flexneri</i>	36	15.2 (33)	-	-	97.1 (34)	-	-	-	8.6 (35)	-
<i>Shigella sonnei</i>	97	66.3 (92)	-	-	95.1 (81)	-	-	-	6.2 (96)	-
<i>Vibrio cholerae</i> O1 Ogawa	34	55.9 (34)	-	-	100 (33)	-	-	-	2.9 (34)	2.9 (34)
<i>Vibrio cholerae</i> non O1, non O139	105	78.1 (105)	-	-	100 (35)	-	-	96 (50)	80 (105)	92 (75)
<i>Vibrio parahaemolyticus</i>	1092	1.9 (1050)	-	96.6 (116)	-	-	-	-	98.9 (1074)	98.2 (952)