

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 45 hospitals , Jan - Dec 2012.

Organism isolated from stool	TOTAL ISOLATES	BETA - LACTAMS	QUINOLONES			AMINOGLYCOSIDE	MISCELLANEOUS			
		AMPICILLIN	NALIDIXIC ACID	CIPROFLOXACIN	NORFLOXACIN	GENTAMICIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> Non-typhoidal	1,039	37.5 (961)	81.8 (44)	98.8 (257)	100 (964)	41.9 (62)	2 (49)	82.3 (237)	83.8 (959)	37.9 (314)
<i>Shigella sonnei</i>	55	74.5 (47)	-		94.4 (36)	-	-	-	17.3 (52)	-
<i>Vibrio cholerae</i> non O1, non O139	115	67 (115)	-	90 (80)	100 (17)	-	-	100 (39)	85.1 (114)	83 (94)
<i>Vibrio parahaemolyticus</i>	795	1.2 (775)	-	93.7 (415)	99.7 (398)	98.6 (73)	-	99.1 (338)	99 (774)	98.4 (693)