

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 46 hospitals , Jan - Dec 2013

Organism isolated from stool	TOTAL ISOLATES	BETA - LACTAMS		QUINOLONES		AMINOGLYC OPEPTIDES	MISCELLANEOUS				
		AMPICILLIN	CEFOXITIN	CIPROFLOXACIN	NORFLOXACIN	GENTAMICIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> Non-typhoidal	777	38.9 (761)	-	94.9 (215)	99.9 (719)	-	-	100 (33)	82.4 (244)	83 (758)	48.9 (1439)
<i>Shigella flexneri</i>	22	18.8 (16)	-	-	95 (20)	-	-	-	-	23.8 (21)	-
<i>Shigella sonnei</i>	67	81.4 (59)	-	-	78.8 (66)	-	-	-	100 (45)	3 (67)	9.1 (44)
<i>Staphylococcus aureus</i>	11	-	88.9 (9)	-	-	100 (7)	85.7 (7)	88.9 (9)	-	100 (8)	-
<i>Vibrio cholerae</i> non O1, non O139	27	88.5 (26)	-	-	100 (17)	-	-	-	-	92.6 (27)	-
<i>Vibrio parahaemolyticus</i>	546	2.4 (495)	-	95.1 (164)	100 (407)	98.6 (71)	-	-	99.6 (263)	98.9 (538)	97.3 (409)