

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 50 hospitals , Jan - Dec 2014

Organism isolated from stool	TOTAL ISOLATES	BETA - LACTAMS		QUINOLONES		GLYCOPEPTIDES	MISCELLANEOUS				
		AMPICILLIN	CEFOXITIN	CIPROFLOXACIN	NORFLOXACIN	GENTAMICIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> Non-typhoidal	786	42.6 (776)	-	64.5 (138)	98.1 (752)	-	-	-	82.7 (196)	81.2 (783)	44.1 (102)
<i>Shigella sonnei</i>	50	72.1 (43)	-	-	75.6 (45)	-	-	-	92.3 (26)	6.2 (48)	5.9 (34)
<i>Vibrio cholerae</i> non O1, non O139	51	82 (50)	-	-	-	-	-	-	-	81.6 (49)	-
<i>Vibrio parahaemolyticus</i>	601	1.2 (511)	-	94.8 (233)	99.7 (338)	100 (82)	-	-	99.4 (172)	98.8 (576)	98.7 (446)