

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 60 hospitals , Jan - Sep 2015

Organism isolated from stool	TOTAL ISOLATES	BETA - LACTAMS		QUINOLONES	GLYCOPEPTIDES	MISCELLANEOUS				
		AMPICILLIN	CEFOXITIN	CIPROFLOXACIN	GENTAMICIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> Non-typhoidal	476	38 (445)	-	68.2 (44)	-	-	-	82.1 (112)	77 (452)	48.4 (95)
<i>Shigella flexneri</i>	12	-	-	-	-	-	-	-	-	-
<i>Shigella sonnei</i>	10	-	-	-	-	-	-	-	-	-
<i>Staphylococcus aureus</i>	4	-	-	-	-	-	-	-	-	-
<i>Vibrio cholerae</i> non O1, non O139	44	65.9 (44)	-	-	-	-	-	-	90.9 (44)	-
<i>Vibrio parahaemolyticus</i>	378	5.3 (360)	-	98.7 (157)	85.7 (28)	-	-	-	97.6 (371)	95.7 (304)