

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 55 hospitals , Jan - Dec 2016

Organism	TOTAL ISOLATES	BETA - LACTAMS												CARBAPENEMS		QUINOLONES					AMINOGLYCOSIDES			GLYCOPEPTIDES		MISCELLANEOUS								
		PENICILLIN	AMPICILLIN	AMOXICILLIN/CLAVULANIC ACID	AMPICILLIN /SULBACTAM	PIPERACILLIN /TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	CIPROFLOXACIN BY MIC	LEVOFLOXACIN	NORFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	NETILMICIN	VANCOMYCIN BY MIC	TEICOPLANIN	CLINDAMYCIN	ERYTHROMYCIN	NITROFURANTOIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Non-typhoidal	744		44.9 (643)															60.2 (196)	96.7 (30)	- ^o	97.4 (546)	- ^o											81.5 (697)	
<i>Shigella</i> spp.	284		69.2 (263)															-	-	-	61 (200)	-											9 (279)	
<i>Vibrio cholerae</i> (all serotypes)	123		65.2 (112)																														-	90.1 (101)
<i>Vibrio cholerae</i> non O1, non O139	91		73.3 (90)																														-	92.4 (79)
<i>Vibrio parahaemolyticus</i>	734		2.8 (580)																														99.1 (233)	98.5 (470)
<i>Vibrio</i> spp.	874		6.7 (670)	90.1 (191)	-	97.1 (172)	- ^o	- ^o	-	93 (215)	97.6 (205)	93.1 (189)	-	-	-	-		97.6 (550)	-	-		99 (205)											98.1 (834)	97.4 (544)