

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 35 hospitals , Jan - Jun 2016

Organism	TOTAL ISOLATES	BETA - LACTAMS														CARBAPENEM		QUINOLONES				AMINOGLYCOSIDES			GLYCOPEPTIDES		MISCELLANEOUS							
		PENICILLIN	AMPICILLIN	AMOXICILLIN/CLAVULANIC ACID	AMPICILLIN/SULBACTAM	PIPERACILIN/TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	OXAICLIN	CEFOXITIN	IMPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	NORFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	METILMICIN	VANCOMYCIN BY E-TEST	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	NITROFURANTOIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE		
<i>Salmonella</i> Non-typhoidal	279	-	47.2 (246)	-	-	-	-	-	-	-	-	-	-	-	-	-	51.8 (56)	- MIC	98.3 (233)	- MIC	-	-	-	-	-	-	-	-	-	-	-	-	78.5 (265)	-
<i>Shigella flexneri</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	- MIC	-	- MIC	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Shigella sonnei</i>	30	-	63.3 (30)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	- MIC	-	- MIC	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Staphylococcus aureus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Vibrio cholerae</i> non O1, non O139	38	-	63.2 (38)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97 (33)	
<i>Vibrio parahaemolyticus</i>	162	-	2 (153)	-	-	-	-	-	-	-	-	-	-	-	-	-	100 (92)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100 (152)	98.2 (111)	