

### Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 74 hospitals, Jan - Dec 2017

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE		
<i>Salmonella</i> , Non-typhoidal	4,896		50.3 (4676)																			84.6 (4725)		
<i>Shigella</i> spp.	107		53.3 (105)											55.3 (47)								20.2 (104)		
<i>Vibrio cholerae</i> (all serotypes)	82		67.1 (76)																			- (76)	80.3 (58)	89.7 (58)
<i>Vibrio cholerae</i> non O1, non O139	70		65.6 (64)																			- (65)	84.6 (65)	94.1 (51)
<i>Vibrio parahaemolyticus</i>	620		2.4 (551)	-	-	-	- <sup>e</sup>	- <sup>e</sup>	91.2 (80)	100 (44)	-	-	-	-	99.5 (371)	-	-	-	97.7 (43)			99.3 (555)	98.4 (448)	
<i>Vibrio</i> spp.	733		7.9 (642)	-	-	-	- <sup>e</sup>	- <sup>e</sup>	93 (100)	100 (54)	-	-	-	-	98.6 (439)	-	-	-	98.1 (52)			98 (656)	97.4 (538)	

