

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 61 hospitals , Jan - Jun 2017

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		MISCELLANEOUS		
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SUL BACTAM	PIPERACILLIN / TAZOBACTAM	CEF AZOLIN (A)	CEF AZOLIN (U)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	CHL ORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Non typhoidal	309		50.7 (274)											68 (103)	- ^e	- ^e					83.3 (288)	
<i>Shigella</i> spp.	38		71.1 (38)																		13.9 (36)	
<i>Vibrio cholerae</i> (all serotypes)	29	-																	-		82.8 (29)	92.6 (27)
<i>Vibrio parahaemolyticus</i>	217			-	-	-	- ^e	- ^e	-	-	-	-	-	100 (147)	-	-	-	-			100 (204)	100 (161)
<i>Vibrio</i> spp.	251			-	-	-	- ^e	- ^e	92.6 (27)	96.3 (27)	-	-	-	-	99.4 (170)	-	-	-	-		100 (228)	100 (193)