

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 44 hospitals, Jan - Sep 2017

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		MISCELLANEOUS			
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE	
<i>Salmonella</i> , Non typhoidal	-		
<i>Shigella</i> spp.	-		
<i>Vibrio cholerae</i> (all serotypes)	35	.																		.	80 (35)	93.9 (33)	
<i>Vibrio parahaemolyticus</i>	178			100 (127)		99.4 (167)	100 (153)
<i>Vibrio</i> spp.	192			100 (131)		99.4 (181)	100 (165)