

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 85 hospitals, Jan - Dec 2018

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS					
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE	
<i>Salmonella</i> , Non-Typhoidal	4701		49.8 (4490)					86 (763)	88.3 (154)	92.1 (1739)						73 (3380)	- <sup>o</sup>									89.4 (982)	86.2 (4472)	
<i>Vibrio cholerae</i> (all serotypes)	81		71.8 (78)																							-	82.3 (79)	90.9 (55)
<i>Vibrio cholerae</i> non O1, non O139	78		70.7 (75)																							-	82.9 (76)	90.6 (53)
<i>Vibrio parahaemolyticus</i>	719		3.6 (631)	-	-	-	- <sup>o</sup>	96.9 (97)	100 (50)		-					98.2 (502)	-	-	98.4 (63)								99.2 (625)	99 (514)
<i>Vibrio spp.</i>	163		45.9 (146)	-	-	-	- <sup>o</sup>	94.4 (36)	-		-					98.4 (128)	-	-	-								84.3 (140)	92.8 (111)

<sup>e</sup>: MIC Interpretive Criteria

