

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 69 hospitals, Jan - Jun 2018

Organism	TOTAL ISOLATES	BETA LACTAMS													CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS						
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN/ SULBACTAM	PIPERACILLIN/ TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	COTRIMOXAZOLE	TETRACYCLINE	
<i>Salmonella</i> , Non-typhoidal	1,633		47.1 (1535)					83.1 (266)	79.1 (43)	94.8 (363)	-						69.4 (1123)	-	-								89.4 (359)	84 (1572)		
<i>Shigella</i> spp.	41		71.1 (38)														36.4 (33)												17.9 (39)	
<i>Vibrio cholerae</i> (all serotypes)	34		80.6 (31)																								-	78.1 (32)	-	
<i>Vibrio cholerae</i> non O1, non O139	34		80.6 (31)																								-	78.1 (32)	-	
<i>Vibrio parahaemolyticus</i>	254		2.7 (220)	-	-	-	-	100 (36)	-		-						99.4 (170)	-	-	-	-							98.2 (222)	98.8 (161)	
<i>Vibrio</i> spp.	44		33.3 (39)	-	-	-	-	-	-		-						-	-	-	-	-						88.6 (35)	93.3 (30)		