

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 47 hospitals, Jan - Sep 2018

Organism	TOTAL ISOLATES	BETA - LACTAMS												CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS					
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE
<i>Salmonella</i> , Non-typhoidal	1,680	-	-	-	-	- <sup>e</sup>	89.5 (238)	94.3 (88)	92.2 (321)	-	-	-	-	-	-	76.4 (1081)	- <sup>e</sup>	- <sup>e</sup>	-	-	-	-	-	-	-	88.9 (395)	85.1 (1605)	-
<i>Vibrio parahaemolyticus</i>	247	4.1 (194)	-	-	-	- <sup>e</sup>	91.5 (59)	-	-	-	-	-	-	-	-	97.5 (157)	-	-	-	100 (41)	-	-	-	-	-	-	97.5 (199)	98.1 (157)
<i>Vibrio</i> spp.	81	28 (75)	-	-	-	- <sup>e</sup>	-	-	-	-	-	-	-	-	-	98.4 (61)	-	-	-	-	-	-	-	-	-	-	91.5 (71)	-

