

Percentage of Susceptible Organisms Isolated From Stool and Rectal Swab, 92 hospitals, Jan - Dec 2019

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Typhoidal	5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Salmonella</i> , Non-Typhoidal	5,381	52.1 (4623)	-	-	-	-	-	-	-	-	-	-	-	-	74.3 (4100)	61.8 [*] (228)	-	-	-	-	-	-	-	-	87.8 (4988)	-	
<i>Shigella</i> spp.	142	42.7 (131)	-	-	-	-	-	-	-	-	-	-	-	-	58.5 (65)	-	-	-	-	-	-	-	-	67.5 (80)	16.4 (140)	-	
<i>Staphylococcus aureus</i>	31	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Vibrio cholerae</i> (all serotypes)	193	76.6 (111)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	88.6 (158)	89.6 (106)	
(O1)	14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
(non O1, non O139)	179	76.8 (99)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91 (144)	90.6 (96)	
<i>Vibrio parahaemolyticus</i>	890	2.5 (637)	-	-	98 (49)	- [*]	-	90.2 (215)	93.2 (117)	90.7 (43)	86 (57)	93.5 (46)	96.4 (56)	59.6 (643)	91.2 (57)	87.1 (31)	97 (67)	-	-	-	-	-	-	-	99.6 (727)	99.1 (552)	
<i>Vibrio</i> spp.	130	56.3 (87)	-	-	-	- [*]	-	90.5 (42)	-	-	-	-	-	-	67 (88)	-	-	-	-	-	-	-	-	-	85.3 (95)	92.9 (85)	

* ข้อมูลความไวจากวิธี disk diffusion และ MIC รวมกัน

