

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 79 hospitals, Jan - Mar 2019

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS					
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE	
<i>Salmonella</i> , Non-Typhoidal	974		46.7 (826)					88.9 (198)	.	93.5 (245)						73 (655)	.									92.7 (262)	86.6 (900)	
<i>Shigella</i> spp.	30			
<i>Vibrio cholerae</i> (all serotypes)	32		.																						.	90.3 (31)	.	
<i>Vibrio cholerae</i> non O1, non O139	30		
<i>Vibrio parahaemolyticus</i>	193		0.8 (126)	89.7 (58)	67.9 (131)	.	.	.								98.7 (156)	100 (99)

