

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 79 hospitals, Jan - Jun 2019

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Non-Typhoidal	2,366		49.9 (1988)					90.4 (481)	90.9 (33)	92.5 (616)						74 (1645)	-								90.6 (596)	87.2 (2185)	
<i>Shigella</i> spp.	75		47 (66)													44.4 (36)	-								61.4 (44)	18.3 (71)	
<i>Vibrio cholerae</i> (all serotypes)	83		72.7 (55)																						-	86.4 (81)	87.5 (40)
<i>Vibrio cholerae</i> non O1, non O139	78		75 (52)																						-	89.6 (77)	86.8 (38)
<i>Vibrio parahaemolyticus</i>	323		0.9 (223)	-	-	-	-	91.7 (96)	94.9 (39)		-	-	-	-	68.6 (223)	-	-	-								99.3 (271)	98.9 (176)

