

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 36 hospitals , Oct - DEC 2013.

Organism isolated from stool	TOTAL ISOLATION	BETA - LACTAM			QUINOLONE			GLYCOPEPTIDE	MISCELLANEOUS				
		AMPICILLIN	OXACILLIN	CEFOXITIN	NALIDIXIC ACID	CIPROFLOXACIN	NORFLOXACIN	GENTAMICIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> Non-typhoidal	113	38.5 (109)	-	-	-	75.8 (33)	100 (100)	-	-	-	85.7 (35)	89.5 (105)	-
<i>Shigella flexneri</i>	7	25 (4)	-	-	-	-	85.7 (7)	-	-	-	75 (4)	14.3 (7)	-
<i>Shigella sonnei</i>	4	-	-	-	-	-	100 (4)	-	-	-	100 (3)	0 (4)	-
<i>Staphylococcus aureus</i>	3	-	-	66.7 (3)	-	-	-	-	66.7 (3)	66.7 (3)	-	-	-
<i>Vibrio cholerae</i> non O1, non O139	10	70 (10)	-	-	-	100 (8)	-	-	-	-	-	90 (10)	57.1 (7)
<i>Vibrio parahaemolyticus</i>	138	1.7 (115)	-	-	-	96.4 (55)	100 (71)	100 (12)	-	-	100 (39)	99.3 (138)	100 (97)