

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, University Hospital, 3 hospitals, JAN - DEC 2017

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		MISCELLANEOUS		
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Non typhoidal	
<i>Shigella</i> spp.	
<i>Vibrio cholerae</i> (all serotypes) □	
<i>Vibrio parahaemolyticus</i>	54			100 (41)		100 (46)	.
<i>Vibrio</i> spp. □	54			100 (41)		100 (46)	.