

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 4 hospitals , Jan - Jun 2017

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE		
<i>Salmonella</i> , Non typhoidal	-				
<i>Shigella</i> spp.	-				
<i>Vibrio cholerae</i> (all serotypes)	-		
<i>Vibrio parahaemolyticus</i>	56			90.9	92.7	.	98.2	.	.		96.3	.			
<i>Vibrio</i> spp.	65			92.1	92.2	.	98.4	.	.		96.8	.			
									(55)					(55)		(55)				(54)				
									(63)					(64)		(64)				(62)				