

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 61 hospitals , Jan - Jun 2017

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		MISCELLANEOUS		
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Non typhoidal	309		50.7 (274)											68 (103)	. ^e	. ^e				83.3 (288)		
<i>Shigella</i> spp.	38		71.1 (38)											.	.	.				13.9 (36)		
<i>Vibrio cholerae</i> (all serotypes)	29	.																	.	82.8 (29)	92.6 (27)	
<i>Vibrio parahaemolyticus</i>	217		 ^e	. ^e	100 (147)		100 (204)	100 (161)	
<i>Vibrio</i> spp.	251			92.6 (27)	96.3 (27)	.	.	.	99.4 (170)		100 (228)	100 (193)	