



Gram negative	<i>Stenotrophomonas maltophilia</i>	510																					72.6									98.7				
	<i>Vibrio cholerae</i> (all serotypes)																						197									374				
	<i>Vibrio cholerae</i> non O1, non O139																																			
	<i>Vibrio parahaemolyticus</i>	32		3.4																			100										100	100		
	<i>Vibrio</i> spp.	45		21.4																			30									29	30			
	<i>Enterococcus faecalis</i>	1,750	78		91.8																		17.7 <sup>a</sup>										62.2	67.4		
	<i>Enterococcus faecium</i>	959	21.8		20.1																		5.2 <sup>u</sup>										95.2	5.1 <sup>f</sup>		
	<i>Enterococcus</i> spp.	191	59.5		66.5																		14.1 <sup>u</sup>										75.5	3.7 <sup>WT</sup>		
	<i>Staphylococcus aureus</i> (all isolates)	2,702	9.5																				88.6 <sup>f</sup>											93.7	52.9	
	(MRSA)	268	0																				0 <sup>f</sup>											74.9	41.4	
	(MSSA)	1,953	11																				100 <sup>f</sup>											98.4	55	
	(ICU)																																			
	(inpatient)	412	0																																	
	(outpatient)	41	10																																	
	<i>Staphylococcus</i> , coagulase negative	3,135	13																																	
	<i>Staphylococcus</i> , coagulase negative (blood)	2,169	55.8																																	
	<i>Streptococcus agalactiae</i>	62	100		100																															
	<i>Streptococcus</i> , β-hemolytic not Group A,B	183	88.9		97.5																															
	<i>Streptococcus pyogenes</i>	563	97.1		99.2																															
	<i>Streptococcus pneumoniae</i> (all isolates)	234	52 <sup>d</sup>																																	
	(age 0-5 years old)	32																																		
	(age > 5 years old)	204	53.7 <sup>d</sup>																																	
	(Sterile sites) <sup>b</sup>	100	66.7 <sup>d</sup>																																	
	(Non-sterile sites) <sup>c</sup>	149	44.8 <sup>d</sup>																																	
	(Meningitis: by E-test)																																			
	(Nonmeningitis: by E-test)																																			

<sup>a</sup>: No CLSI Interpretive Criteria. Interpret according to cefoperazone/sulbactam in *Enterobacteriaceae*

<sup>b</sup>: Blood, Pleural Fluid

<sup>c</sup>: Sputum, Ear, Sinus

<sup>d</sup>: Interpret according to oxacillin susceptibility test

<sup>e</sup>: MIC Interpretive Criteria

<sup>f</sup>: Interpret according to ceftoxitin susceptibility test

<sup>h</sup>: High-Level Aminoglycoside

<sup>u</sup>: Urine, Urine Catheter, Urine Clean- Voided

<sup>WT</sup>: Wild-type