





Percentage of susceptible Organisms Isolated From All Specimen, 8 hospitals, RMSc 11&11/1, Jan - Jun 2017

Organism		TOTAL ISOLATES	BETA - LACTAMS														CARBAPENEMS			POLY MYXINS	QUINOLONES				AMINOGLYCOSIDES			GLYCOPEPTIDES			MISCELLANEOUS														
			PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFUROXIME SODIUM (parenteral)	CEFUROXIME SODIUM (Oral)	CEFOPERAZONE / SULBACTAM	CEFTOXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILLIN	CEFOXITIN	BERTAPENEM	IMPENEM	MEROPENEM	COLISTIN BY MIC	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	NORFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	GENTAMICIN 120 µg	NETILMICIN	VANCOMYCIN	VANCOMYCIN BY MIC	TEICoplanin	FOSFOMYCIN	CLINDAMYCIN	ERYTHROMYCIN	NITROFURANTOIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE				
Gram positive	(Sterile sites) <sup>b</sup>	63	77.1 (39)	-	-	-	-	-	-	-	-	-	-	-	-	-	87.1 (35)	30.6 (36)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	(Non-sterile sites) <sup>c</sup>	134	67.8 (59)	-	100 (42)	-	-	-	-	-	-	-	-	-	-	-	67.8 (59)	-	-	-	-	-	100 (83)	-	100 (35)	-	-	-	-	-	100 (107)	-	-	-	-	-	-	78.9 (38)	80.6 (36)	-	-	90 (30)	-	-	-
	(Meningitis: by E-test)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	(Nonmeningitis: by E-test)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

<sup>a</sup> : No CLSI Interpretive Criteria. Interpret according to ceftoperazone/sulbactam in *Enterobacteriaceae*

<sup>b</sup> : Blood, Pleural Fluid

<sup>c</sup> : Sputum, Ear, Sinus

<sup>d</sup> : Interpret according to oxacillin susceptibility test

<sup>e</sup> : MIC Interpretive Criteria

<sup>f</sup> : Interpret according to cefoxitin susceptibility test

<sup>h</sup> : High-Level Aminoglycoside

<sup>u</sup> : Urine, Urine Catheter, Urine Clean- Voided

<sup>WT</sup> : Wild-type