

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 4 hospitals, BKK, Jan - Dec 2016

Organism	TOTAL ISOLATES	BETA - LACTAMS												CARBAPENEM		QUINOLONES					AMINOGLYCOSIDES			GLYCOPEPTIDES		MISCELLANEOUS						
		PENICILLIN	AMPCILLIN	AMOXICILIN/CLAVULANIC ACID	AMPCILLIN/SULBACTAM	PIPERACILLIN/TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFUROXIME SODIUM (parenteral)	CEFTAZIME	CEFTAZIDIME	CEFERPIME	OXAGLIN	CEFOXITIN	IMIPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	NORFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	NETILMICIN	VANCOMYCIN BY E-TEST	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	NITROFURANTON	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella, typhoidal</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Shigella</i> spp.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Staphylococcus aureus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Vibrio cholerae</i> (all serotypes)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Vibrio cholerae</i> O1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Vibrio cholerae</i> O139	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Vibrio cholerae</i> non O1, non O139	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Vibrio</i> spp.	71	-	4.8 (62)	-	-	-	-	e	e	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96.7 (60)	100 (61)