

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, RMsC 1 & 1/1, 11 hospitals, Jan - Jun 2018

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS						
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE
<i>Salmonella</i> , Typhoidal	-	-	-					-	-	-	-						-	- <sup>e</sup>	- <sup>e</sup>							-	-	
<i>Salmonella</i> , Non-typhoidal	248		34.8 (210)					85.4 (48)	-	-	-						74.2 (93)	- <sup>e</sup>	- <sup>e</sup>							77.9 (86)	80.2 (242)	
<i>Shigella</i> spp.	-	-	-													-	-	-	-									-
<i>Staphylococcus aureus</i>	-	-										-	f				-	-	-		-	- <sup>e</sup>	-	-	-	-	-	-
<i>Vibrio cholerae</i> (all serotypes)	-	-	-																							-	-	-
<i>Vibrio cholerae</i> O1	-	-	-																							-	-	-
<i>Vibrio cholerae</i> non O1, non O139	-	-	-																							-	-	-
<i>Vibrio parahaemolyticus</i>	-	-	-	-	-	- <sup>e</sup>		-	-		-	-	-	-		-	-	-	-	-	-					-	-	-
<i>Vibrio</i> spp.□	-	-	-	-	-	- <sup>e</sup>		-	-	-	-	-	-	-		-	-	-	-	-						-	-	