Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, RMSc3, 5 hospitals, January -June 2018

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES				AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMIPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICOPLANIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
Salmonella, Typhoidal	-																		e	e									
Salmonella, Non-typhoidal	81		(51.6) (62)															(92.5) (67)	e	e								93.9 66	
Shigella spp.	-																												
Staphylococcus aureus	-												f											e					
Vibrio cholerae (all serotypes)	-																												
Vibrio cholerae O1	-																												
Vibrio cholerae non O1, non O139	-																												
Vibrio parahaemolyticus	-						e																						
Vibrio spp.	-						e																						