Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 8 hospitals, JANUARY - JUNE 2018, RMSC 5th

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES				MINOGLYCOSIDE		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/CLAVULANIC ACID	AMPICILLIN /SULBACTAM	PIPERACILLIN /TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMIPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICOPLANIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
Salmonella, Typhoidal	-																		e	e									
Salmonella, Non-typhoidal	340		45.4 (337)						77.6 (134)									51.4 (286)	e	e								83.2 (339)	
Shigella spp.	5																												
Staphylococcus aureus	-												f											e					
Vibrio cholerae (all serotypes)	30																												
Vibrio cholerae O1	-																												
Vibrio cholerae non O1, non O139	30																												
Vibrio parahaemolyticus	76		1.6 (63)				e											100 (73)										98.5 (65)	100 (52)
Vibrio spp.	5						e																						