

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 7 hospitals,RMSc 8, jan -Jun 2018

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE
<i>Salmonella</i> , Typhoidal	-		-					-	-	-	-						-	- ^e	- ^e							-	-	
<i>Salmonella</i> , Non-typhoidal	87		54.8 (84)					-	-	-	-						62.8 (86)	- ^e	- ^e							90.7 (54)	86.5 (74)	
<i>Shigella</i> spp.	-		-													-	-	-									-	
<i>Staphylococcus aureus</i>	-	-										- ^f					-	-	-		-	-	- ^e	-	-	-	-	-
<i>Vibrio cholerae</i> (all serotypes)	-		-																							-	-	-
<i>Vibrio cholerae</i> O1	-		-																							-	-	-
<i>Vibrio cholerae</i> non O1, non O139	-		-																							-	-	-
<i>Vibrio parahaemolyticus</i>	-		-	-	-	-	- ^e	-	-		-	-	-	-	-		-	-	-	-	-					-	-	
<i>Vibrio</i> spp.	-		-	-	-	-	- ^e	-	-		-	-	-	-	-		-	-	-	-	-					-	-	

%S				
<50	50-69	70-79	80-89	≥90