

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, RMSC 7, 4 hospitals, Jan - Jun 2018

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES				AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS					
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parentera)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE
<i>Salmonella</i> , Typhoidal	-		
<i>Salmonella</i> , Non-typhoidal	-		
<i>Shigella</i> spp.	-		
<i>Staphylococcus aureus</i>	-
<i>Vibrio cholerae</i> (all serotypes)	-		
<i>Vibrio cholerae</i> O1	-		
<i>Vibrio cholerae</i> non O1, non O139	-		
<i>Vibrio parahaemolyticus</i>	-	
<i>Vibrio</i> spp. □	-	