

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 65 hospitals , Jan - Dec 2015

Organism isolated from stool	TOTAL ISOLATES	BETA - LACTAMS		QUINO LONES	GLYCO PEPTIDES	MISCELLANEOUS				
		AMPICILLIN	CEFOXITIN	CIPROFLOXACIN	GENTAMICIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> Non-typhoidal	739	40.2 (694)	-	65.5 (116)	-	-	-	81.2 (202)	79.1 (707)	56.3 (158)
<i>Shigella flexneri</i>	-	-	-	-	-	-	-	-	-	-
<i>Shigella sonnei</i>	-	-	-	-	-	-	-	-	-	-
<i>Staphylococcus aureus</i>	-	-	-	-	-	-	-	-	-	-
<i>Vibrio cholerae</i> non O1, non O139	57	61.4 (57)	-	-	-	-	-	-	89.5 (57)	-
<i>Vibrio parahaemolyticus</i>	467	4.2 (432)	-	98.1 (216)	83.7 (43)	-	-	-	98.4 (447)	96.6 (357)