

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 49 hospitals, Jan - Sep 2019

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Typhoidal	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	- ^e	-	-	-	-	-	-	-	-	-	
<i>Salmonella</i> , Non-Typhoidal	2,419	-	53 (2066)	-	-	-	94.1 (303)	-	95.8 (594)	-	-	-	-	-	-	78.5 (1768)	- ^e	-	-	-	-	-	91.3 (713)	88.4 (2245)	-	-	
<i>Shigella</i> spp.	99	-	44 (91)	-	-	-	-	-	-	-	-	-	-	-	-	43.3 (30)	-	-	-	-	-	-	72.3 (65)	12.4 (97)	-	-	
<i>Staphylococcus aureus</i>	30	-	-	-	-	-	-	-	-	-	- ^f	-	-	-	-	-	-	-	-	- ^e	-	-	-	-	-		
<i>Vibrio cholerae</i> (all serotypes)	76	-	76.1 (71)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	85.9 (71)	92.2 (51)		
<i>Vibrio cholerae</i> O1	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
<i>Vibrio cholerae</i> non O1, non O139	72	-	77.9 (68)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	88.2 (68)	91.8 (49)		
<i>Vibrio parahaemolyticus</i>	387	-	2.1 (281)	-	-	- ^e	93.7 (79)	98 (51)	-	-	-	-	-	-	64.3 (258)	-	-	-	-	-	-	-	-	99.3 (304)	98.2 (226)		
<i>Vibrio</i> spp.	45	-	-	-	-	- ^e	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

