

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 83 hospitals, Jan - Dec 2020

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS						
		PENICILLIN	AMPICILLIN	AMOXICILLIN / CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Typhoidal	4		66.7 (3)					50.0 (2)	100.0 (2)	100.0 (1)					25.0 (4)	0.0 e (2)									-	75.0 (4)	
<i>Salmonella</i> , Non-typhoidal	2606		60.5 (2606)					92.5 (919)	92.2 (425)	94.2 (1057)						76.0 (2455)	62.8 e (282)								89.3 (363)	89.6 (2601)	
<i>Shigella</i> spp.	61		45.9 (61)													62.1 (29)	100.0 (6)								77.3 (44)	28.9 (38)	
<i>Staphylococcus aureus</i>	16	0.0 (13)										93.8 f (16)				100.0 (2)	100.0 (1)		100.0 (6)	-	- e	85.7 (7)	85.7 (7)	-	100.0 (6)	0.0 (1)	
<i>Vibrio cholerae</i> (all serotypes)	113		81.6 (103)																					100.0 (7)	93.8 (113)	94.8 (77)	
<i>Vibrio cholerae</i> O1	3		33.3 (3)																					-	66.7 (3)	50.0 (2)	
<i>Vibrio cholerae</i> non O1, non O139	110		83.0 (100)																					100.0 (7)	94.5 (110)	96.0 (75)	
<i>Vibrio parahaemolyticus</i>	358		2.0 (297)	93.2 (59)	87.5 (8)	100.0 (51)	- e	-	94.1 (205)	97.5 (121)		92.0 (50)	94.9 (39)	100.0 (51)	100.0 (64)	79.3 (358)	95.8 (71)	96.1 (51)	96.0 (75)						98.3 (352)	99.6 (234)	
<i>Vibrio</i> spp.	61		31.5 (54)	84.0 (25)	100.0 (1)	100.0 (25)	- e	-	90.9 (44)	100.0 (31)		100.0 (25)	80.0 (25)	64.0 (25)	96.0 (25)	77.0 (61)	92.3 (26)	100.0 (24)	96.4 (28)						91.7 (60)	83.3 (18)	

