

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 79 hospitals, Jan - Mar 2020

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , typhoidal	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Salmonella</i> , Non-typhoidal	909	-	52.6 (797)	-	-	-	-	-	-	-	-	-	-	-	-	74.9 (732)	84 <sup>a</sup> (50)	-	-	-	-	-	-	-	87.4 (874)	-	
<i>Shigella</i> spp.	25	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Staphylococcus aureus</i>	15	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Vibrio cholerae</i> (all serotypes)	38	-	82.9 (35)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	89.5 (38)	-		
(O1)	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
(non O1, non O139)	36	-	84.8 (33)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	88.9 (36)	-		
<i>Vibrio parahaemolyticus</i>	79	-	1.6 (63)	-	-	-	- <sup>a</sup>	-	-	-	-	-	-	-	-	86.4 (59)	-	-	-	-	-	-	-	98.6 (70)	100 (53)		
<i>Vibrio</i> spp.	7	-	-	-	-	-	- <sup>a</sup>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

<sup>a</sup> % susceptible organisms by disk diffusion and MIC method

