

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 79 hospitals, Jan - Jun 2020

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , typhoidal	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	- <sup>e</sup>	-	-	-	-	-	-	-	-	-	
<i>Salmonella</i> , Non-typhoidal	2,093	-	58.6 (1829)	-	-	-	-	-	-	-	-	-	-	-	-	76.3 (1771)	71.2 <sup>e</sup> (198)	-	-	-	-	-	-	-	88.7 (1967)	-	
<i>Shigella</i> spp.	47	-	37.8 (45)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	11.1 (45)	-	
<i>Staphylococcus aureus</i>	26	-	-	-	-	-	-	-	-	-	- <sup>f</sup>	-	-	-	-	-	-	-	-	- <sup>e</sup>	-	-	-	-	-	-	
<i>Vibrio cholerae</i> (all serotypes)	99	-	80.4 (92)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	93.9 (98)	95.3 (64)	
(O1)	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
(non O1, non O139)	96	-	82 (89)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94.7 (95)	96.8 (63)	
<i>Vibrio parahaemolyticus</i>	247	-	1 (193)	-	-	-	- <sup>e</sup>	-	93.7 (79)	95.9 (49)	-	-	-	-	-	73.5 (196)	-	-	97.6 (41)	-	-	-	-	-	99.5 (214)	99.3 (148)	
<i>Vibrio</i> spp.	27	-	-	-	-	- <sup>e</sup>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

\*נתונים מבוססים על דיוקן דיסקים ו-MIC

