

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 87 hospitals, Jan - Dec 2021

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN / CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Typhoidal	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Salmonella</i> , Non-typhoidal	3448	-	54.1 (2966)	-	-	-	-	-	-	-	-	-	-	-	-	73.2 (3205)	62.5 * (275)	-	-	-	-	-	-	-	-	86 (3240)	
<i>Shigella</i> spp.	41	-	78.3 (23)	-	-	-	-	-	-	-	-	-	-	-	-	63.2 (38)	-	-	-	-	-	-	-	-	-	61.1 (36)	
<i>Staphylococcus aureus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Vibrio cholerae</i> (all serotypes)	89	-	69.6 (79)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	81.7 (82)	98.4 (62)
(O1)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
(non O1, non O139)	84	-	70.3 (74)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	80.8 (78)	98.3 (60)
<i>Vibrio parahaemolyticus</i>	442	-	3.1 (290)	97 (67)	-	98.3 (58)	-	-	94.2 (208)	99.3 (145)	-	88.9 (54)	-	95.5 (44)	98 (50)	96.6 (59)	75.3 (365)	93.2 (59)	94.4 (72)	98 (100)	-	-	-	-	-	98.0 (354)	98.7 (236)
<i>Vibrio</i> spp.	70	-	37.8 (45)	-	-	-	-	-	89.1 (46)	90.7 (43)	-	90.9 (22)	-	-	-	83.1 (65)	-	-	-	-	-	-	-	-	-	90.7 (54)	-

* ข้อมูลจากวิธี disk diffusion และ MIC รวมกัน

