

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 94 hospitals, Jan - Dec 2024

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN / CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
Salmonella, Typhoidal	16		-														-	- e								-	
Salmonella, Non-typhoidal	4123		60.5 (3625)														71.8 (4014)	- e								90.2 (3988)	
Shigella spp.	64		34.6 (52)														33.3 (63)	-								14.8 (61)	
Staphylococcus aureus	2	-										- f					-	-		-	-	- e	-	-	-	-	-
Vibrio cholerae (all serotypes)	113		66.3 (101)																						-	86.8 (106)	97.4 (77)
(O1)	6		-																						-	-	-
(non O1, non O139)	104		67.0 (94)																						-	85.9 (99)	97.3 (75)
Vibrio parahaemolyticus	336		10.3 (232)	100 (78)	-	98.6 (74)	- e	-	95.6 (90)	97.2 (108)		98.4 (64)		-	98.7 (77)	98.7 (78)	99.4 (325)	100 (55)	95.0 (80)	98.5 (67)						96.8 (281)	99.4 (164)
Vibrio spp.	53		48.5 (33)	-	-	-	- e	-	-	-		-		-	-	-	96.1 (51)	-	-	-						80.0 (45)	-

*N <30 isolates tested. The result is not significant as per CLSI

* ข้อมูลความไวจากวิธี disk diffusion และ MIC รวมกัน

