

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 107 hospitals, Jan - Dec 2025

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN / CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
Salmonella, Typhoidal	24		-														-	- e								-	
Salmonella, Non-typhoidal	3817		63.4 (3428)														69.8 (3649)	- e								89.8 (3731)	
Shigella spp.	24		-														-	-								-	
Staphylococcus aureus	6	-										- f					-	-		-	-	- e	-	-	-	-	-
Vibrio cholerae (all serotypes)	100		73.7 (95)																						-	89.7 (97)	98.4 (62)
(O1)	7		-																						-	-	-
(non O1, non O139)	86		75.0 (84)																						-	90.5 (84)	98.3 (59)
Vibrio parahaemolyticus	328		11.5 (243)	95.8 (143)	85.2 (54)	100.0 (128)	- e	-	95.3 (106)	95.9 (148)		94.5 (73)		-	100.0 (138)	98.6 (141)	98.8 (320)	99.1 (116)	97.9 (143)	96.0 (99)						95.1 (284)	94.9 (138)
Vibrio spp.	47		45.5 (33)	-	-	-	- e	-	90.3 (31)	94.1 (34)		-		-	-	-	89.1 (46)	96.7 (30)	-	-						85.7 (42)	-

*N <30 isolates tested. The result is not significant as per CLSI guidelines

* ข้อมูลความไวจากวิธี disk diffusion และ MIC รวมกัน

