

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 40 hospitals , Jan - Sep 2014

Organism isolated from stool	TOTAL ISOLATES	BETA - LACTAMS		QUINOLONES		Aminoglyco sides	MISCELLANEOUS				
		AMPICILLIN	CEFOXITIN	CIPROFLOXACIN	NORFLOXACIN	GENTAMICIN	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> Non-typhoidal	605	42.6 (601)	-	62.8 (86)	98.6 (585)	-	-	-	83.5 (170)	80.3 (603)	45.5 (88)
<i>Shigella sonnei</i>	44	73 (37)	-	-	81.6 (38)	-	-	-	96 (25)	7 (43)	6.7 (30)
<i>Vibrio cholerae</i> non O1, non O139	36	80.6 (36)	-	-	100 (2)	-	-	-	-	80 (35)	-
<i>Vibrio parahaemolyticus</i>	417	1.6 (370)	-	96 (176)	100 (228)	100 (51)	-	-	99.2 (119)	98.8 (403)	98.4 (304)