

Percent Susceptibility of Organisms Isolated From Stool and Rectal Swab, 8 hospitals , Jan - Dec 2016, RMSCs 5th.

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEM		QUINOLONES					AMINOGLYCOSIDES			GLYCOPEPTIDES			MISCELLANEOUS									
		PENICILLIN	AMPICILLIN	AMOXICILLIN/CLAVULANIC ACID	AMPICILLIN/SULBACTAM	PIPERACILLIN/TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	OXACILIN	CEFOXITIN	IMIPENEM	MEROPENEM	NALIDIXIC ACID	CIPROFLOXACIN	LEVOFLOXACIN	NORFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	NETILMICIN	VANCOMYCIN BY E-TEST	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	NITROFURANTOIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE			
<i>Salmonella</i> Non-typhoidal	-																																		
<i>Shigella</i> spp.	21																																		
<i>Shigella sonnei</i>	-																																		
<i>Staphylococcus aureus</i>	-																																		
<i>Vibrio cholerae</i> Vibrio cholerae non O1, non O139	66		68.2 (66)																															95.5 (66)	
<i>Vibrio parahaemolyticus</i>	-																																		
<i>Vibrio</i> spp.	230		6.1 (229)																															99.6 (229)	99.5 (219)