

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, RMsC 12 & 12/1, 8 hospitals, Jan -Dec 2017

Organism	TOTAL ISOLATES	BETA - LACTAMS										CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		MISCELLANEOUS			
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFAZOLIN (U)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Non typhoidal	36		58.3 (36)											.	. ^e	. ^e				94.3 (35)		
<i>Shigella</i> spp. ^e	. ^e				.		
<i>Vibrio cholerae</i> (all serotypes)	
<i>Vibrio parahaemolyticus</i>	66		 ^e	. ^e		98.4 (62)	98.3 (58)
<i>Vibrio</i> spp.	73		 ^e	. ^e		98.6 (69)	96.9 (64)