

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 6 hospitals, Jan - Dec 2024 , เขตสุขภาพที่ 13

Organism	TOTAL ISOLATES	BETA - LACTAMS														CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN / CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE	
<i>Salmonella</i> , Typhoidal	2		-																									
<i>Salmonella</i> , Non-typhoidal	374		47.2 (373)													44.4 (374)	- °									85.8 (373)		
<i>Shigella</i> spp.			-													-	-								-			
<i>Staphylococcus aureus</i>	1	-										- f				-	-		-	-	- °	-	-	-	-	-		
<i>Vibrio cholerae</i> (all serotypes)	10		-																						-	-		
(O1)			-																						-	-		
(non O1, non O139)	10		-																						-	-		
<i>Vibrio parahaemolyticus</i>	11		-	-	-	-	- °	-	-	-		-	-	-	-	-	-	-	-						-	-		
<i>Vibrio</i> spp.	4		-	-	-	-	- °	-	-		-	-	-	-	-	-	-	-	-						-	-		

*N <30 isolates tested. The result is not significant as per CLSI g1

* ข้อมูลความไวจากวิธี disk diffusion และ MIC รวมกัน

